

A Dark Hydrogen Cloud in the Virgo Cluster

Robert Minchin¹, Jonathan Davies¹, Michael Disney¹, Peter Boyce², Diego Garcia¹,
Christine Jordan³, Virginia Kilborn⁴, Robert Lang¹, Sarah Roberts¹, Sabina Sabatini⁵

and

Wim van Driel⁶

ABSTRACT

VIRGOHI21 is an HI source detected in the Virgo Cluster survey of Davies et al. (2004) which has a neutral hydrogen mass of $10^8 M_\odot$ and a velocity width of $\Delta V_{20} = 220 \text{ km s}^{-1}$. From the Tully-Fisher relation, a galaxy with this velocity width would be expected to be 12th magnitude or brighter; however deep CCD imaging has failed to turn up a counterpart down to a surface-brightness level of $27.5 \text{ B mag arcsec}^{-2}$. The HI observations show that it is extended over at least 16 kpc which, if the system is bound, gives it a minimum dynamical mass of $\sim 10^{11} M_\odot$ and a mass to light ratio of $M_{\text{dyn}}/L_B > 500 M_\odot/L_\odot$. If it is tidal debris then the putative parents have vanished; the remaining viable explanation is that VIRGOHI21 is a dark halo that does not contain the expected bright galaxy. This object was found because of the low column density limit of our survey, a limit much lower than that achieved by all-sky surveys such as HIPASS. Further such sensitive surveys might turn up a significant number of the dark matter halos predicted by Dark Matter models.

¹School of Physics and Astronomy, Cardiff University, Cardiff, CF24 3YB, UK; Robert.Minchin@astro.cf.ac.uk, Jonathan.Davies@astro.cf.ac.uk, Mike.Disney@astro.cf.ac.uk, Diego.Garcia@astro.cf.ac.uk, LangRH@cardiff.ac.uk, Sarah.Roberts@astro.cf.ac.uk

²Planning Division, Cardiff University, Park Place, Cardiff, CF10 3UA, UK; BoyceP@cardiff.ac.uk

³Jodrell Bank Observatory, University of Manchester, Macclesfield, Cheshire, SK11 9DL, UK; caj@jb.man.ac.uk

⁴Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Mail 31, P.O. Box 218, Hawthorn, Victoria 3122, Australia; vkilborn@astro.swin.edu.au

⁵Osservatorio Astronomico di Roma, via Frascati 33, I-00040, Monte Porzio, Italy; sabatini@mporzio.astro.it

⁶Observatoire de Paris, GEPI, CNRS UMR 8111 and Université Paris 7, 5 place Jules Janssen, F-92195 Meudon Cedex, France; Wim.vanDriel@obspm.fr

Subject headings: dark matter — galaxies: general — galaxies: clusters: individual (Virgo) — radio lines: galaxies

1. Introduction

Simulations of Cold Dark Matter (CDM) models predict far more dark matter halos than are observed as galaxies (Klypin et al. 1999; Moore et al. 1999). For this reason, it has been hypothesized that there must exist dark matter halos that contain no stars (e.g. Jimenez et al. 1997; Verde et al. 2002). The advent of neutral hydrogen multibeam systems has allowed surveys of large areas of sky to be carried out with much higher sensitivity than has been possible in the past, thus allowing sources to be detected by their gas content alone rather than their stars and opening up the possibility of finding truly isolated clouds of extragalactic gas with no stars. Prior to this, blind HI surveys either covered very small areas or were insensitive to HI column densities lower than $\sim 10^{20} \text{ cm}^{-2}$ ($\sim 1M_{\odot} \text{ pc}^{-2}$) (Minchin et al. 2003).

Davies et al. (2004) used the multibeam system on the Lovell telescope at Jodrell Bank Observatory to carry out a deep neutral-hydrogen (HI) survey of the Virgo Cluster (VIRGOHI), covering 32 square degrees and detecting 31 sources. Of these sources, 27 were known cluster members and 4 were new detections. One of the latter lay behind M86 and was thus unobservable optically and one was undetected in follow-up observations and is therefore believed to be a false detection. The other two were confirmed at Arecibo and flagged by Davies et al. as possible isolated HI clouds. One (VIRGOHI27) has an optical counterpart visible in our deep CCD images, the other (VIRGOHI21, the subject of this letter) does not.

There have been several previous claims of the detection of isolated clouds of extragalactic gas with no stars in them, but subsequent analyses have either revealed the optical counterparts (Giovanelli & Haynes 1989; McMahon et al. 1990), or shown that the gas is merely debris from nearby visible galaxies (Schneider et al. 1983). Many other detections of HI clouds have been associated with nearby optically-bright galaxies (Kilborn et al. 2000; Boyce et al. 2001; Ryder et al. 2001). VIRGOHI21 cannot be so easily explained.

2. Further observations

Following detection, VIRGOHI21 was re-observed at Arecibo. The observations are fully described by Davies et al. (2004), here we give a much more detailed analysis of the

Arecibo data and present new VLA and optical observations. The Arecibo observations used a number of pointings in a pattern around the best-fit location from the Jodrell Bank data, leading to the source being detected in five of the Arecibo beams.

Fig. 1 shows the single-dish spectra of VIRGOHI21. Spectrum (a) is the discovery spectrum from Jodrell Bank, this has a noise level of 4 mJy per 13.2 km s^{-1} channel and a 5σ column-density sensitivity $N_{\text{HI},\text{lim}} = 7 \times 10^{18} \text{ cm}^{-2}$ if spread over 200 km s^{-1} . From it we measure a total flux of $F_{\text{HI}} = 2.4 \pm 0.3 \text{ Jy km s}^{-1}$ and a velocity width at 20 per cent of the peak flux of $\Delta V_{20} = 290 \text{ km s}^{-1}$. Spectra (b), (c) and (d) are three north through south beams across the source from the Arecibo observations (labelled (b) - (d) in Fig. 2). These have a noise level of 1.3 mJy per 5.5 km s^{-1} channel, giving $N_{\text{HI},\text{lim}} = 2.7 \times 10^{19} \text{ cm}^{-2}$ over 200 km s^{-1} . They reveal a systematic increase in velocity of $\sim 200 \text{ km s}^{-1}$ from south to north. Spectrum (e) is the co-added spectrum from all 16 Arecibo beams (shown in Fig. 2) from which we measure $F_{\text{HI}} = 3.8 \pm 0.2 \text{ Jy km s}^{-1}$ and $\Delta V_{20} = 220 \text{ km s}^{-1}$.

Fig. 2 shows the Arecibo pointing pattern for VIRGOHI21 and which beams made firm detections (better than 4σ). It can be seen that VIRGOHI21 is extended over at least one Arecibo beam width $\approx 3.6 \text{ arc min}$, or 16 kpc at an assumed distance to Virgo of 16 Mpc (Graham et al. 1999). Using the measured HI flux, we calculate an HI mass of $2 \times 10^8 M_{\odot}$ if it is at the distance of the Virgo Cluster, or $7 \times 10^8 M_{\odot}$ if it is at its Hubble distance (29 Mpc for $H_0 = 70 \text{ Mpc}^{-1} \text{ km s}^{-1}$). For the rest of this letter we will assume the former as more conservative in this context. The best-position for the center of the HI emission, formed by weighting the Arecibo detection positions by their fluxes, is $12^{\text{h}}17^{\text{m}}53.6^{\text{s}}, +14^{\circ}45'25''$ (J2000). We can dismiss the possibility that this is side-lobe emission from another part of the sky, because VIRGOHI21 has been detected with two telescopes with very different side-lobes. Additionally, there are no HI-massive galaxies in the region that match its velocity profile (Davies et al. 2004).

HI observations with the VLA in D-array in August 2004 reached a 5σ column-density limit of 10^{20} cm^{-2} over 60 km s^{-1} in 6 hours (0.5 mJy per 20 km s^{-1} channel with a beam size of $48 \times 45 \text{ arc seconds}$ using natural weighting). Solar interference on the shorter baselines meant that the observations did not reach the hoped-for column-density sensitivity; most of the data from baselines shorter than $\sim 270 \text{ m}$ ($\sim 1.3\text{k}\lambda$) had to be flagged as bad. These observations did detect compact HI associated with a nearby dwarf elliptical (MAPS-NGP O_435_1291894) at a different velocity, but, although sensitive to compact, narrow-line gas down to a 5σ limit of $3 \times 10^6 M_{\odot}$, were not sensitive enough to low column-density, high velocity-width gas to detect VIRGOHI21. For the Arecibo and VLA observations to be consistent, the source must again exceed 16 kpc in diameter.

We have obtained deep optical CCD images in B , r and i bands with the 2.5-m Isaac

Newton Telescope (INT). By binning the B -band image (generally the best band for looking for low surface brightness galaxies) into 1 arc sec pixels, we reach a surface-brightness limit of $27.5 B \text{ mag arcsec}^{-2}$; treating the r and i images in the same way gives surface-brightness limits of ~ 27.0 and of $25.8 \text{ mag arcsec}^{-2}$ respectively. Previous experience indicates that, on the B -band frame, we should be able to easily detect objects of ten arc sec scale or larger at this surface brightness limit (Sabatini et al. 2003; Roberts et al. 2004). This is more than 100 times dimmer than the central surface brightness of the disks of typical spiral galaxies ($21.5 B \text{ mag arcsec}^{-2}$, Freeman 1970) and dimmer than any known massive low surface-brightness galaxy ($26.5 B \text{ mag arcsec}^{-2}$, Bothun et al. 1987) or (for typical $B - V$ colors of ~ 0.6) than the lowest surface-brightness dwarf galaxy ($26.8 V \text{ mag arcsec}^{-2}$ Zucker et al. 2004).

Although we easily were able to identify an optical counterpart (at $12^h26^m40.1^s, +19^\circ45'50''$ J2000) to the other possible H I cloud, VIRGOHI27, no optical counterpart to VIRGOHI21 is visible down to our surface-brightness limit on any of our deep images of this region (Fig. 3), nor can one be found with advanced routines for detecting low surface-brightness galaxies (matched filtering and wavelets, Sabatini et al. 2003). Unlike VIRGOHI27, the bluest objects in the field, which might be associated with H II regions, are widely distributed without any concentration towards the H I center. Looking at the statistics of the sky noise for the frame, the mean number of counts and the standard deviation (excluding stars) are similar in the area of the H I detection to other, blank areas of sky in the vicinity. The number of detected faint objects is not significantly above the average in a box centred on the H I position: there are 3 objects with $m_B > 23$ in a 100×100 pixel (33.3×33.3 arc sec) region centred on our best H I position, compared to an average of 1.8 ± 1.4 across the cube, and 4 objects with $m_B > 22$ compared to an average of 2.6 ± 1.7 .

As is to be expected there are some features on the image of VIRGOHI21 that are obviously faint galaxies. These are labelled (A) - (E) in Fig. 3. (A) is a small source with a star superposed (which prevents accurate determination of its color and luminosity) 2 arc minutes north of the weighted center, just within the Full Width Half Maximum (FWHM) of the strongest H I beam. Its east-west orientation is at odds with the north-south orientation expected from the velocity field of VIRGOHI21 and, if VIRGOHI21 is rotating, this galaxy is too far north for the rotation to be centred on its position. Neither its size, position nor orientation make this a likely optical counterpart to VIRGOHI21. (B) is another uncatalogued galaxy, 3.5 arc minutes south-west of the weighted center. It lies within the FWHM of an Arecibo beam where there was no detection, thus it cannot be the optical counterpart. There are three objects classified as galaxies within six arc minutes. One (C) at the very edge of one of the Arecibo beams is a dwarf elliptical (MAPS-NGP O_435_1291894) that is detected in our VLA data at 1750 km s^{-1} (see Fig. 1). The H I is separated both spatially and in velocity from VIRGOHI21 and so it cannot therefore be the

optical counterpart. Another of the catalogued galaxies (D - MAPS-NGP O_435_1292289) is a double star miscatalogued as a galaxy, and the third (E - VCC 0273) lies in an Arecibo beam where no detection was made.

We conclude that there is no optical counterpart to VIRGOHI21 down to a B -band surface-brightness limit of $27.5 B \text{ mag arcsec}^{-2}$. This is less than 1 solar luminosity pc^{-2} , giving a maximum luminosity in stars of $\sim 10^8$ solar luminosities if a diameter of 16 kpc is assumed. If VIRGOHI21 is a bound system (see discussion below), this leads to a mass to light ratio in solar units of $M_{\text{dyn}}/L_B > 500$ compared to a typical L^* galaxy like the Milky Way with $M_{\text{dyn}}/L_B \sim 50$ within its HI radius (Salucci & Persic 1997). For standard stellar M/L_B ratios the upper limit on the mass in stars is approximately equal to the mass in HI.

3. Discussion

The closest bright HI-rich galaxies ($M_B < -16$, $M_{HI} > 10^8 M_\odot$) are shown in Fig. 2. These are NGC 4262 at 1489 km s^{-1} and NGC 4254 at 2398 km s^{-1} , each at projected distances of 120 kpc away to the east and southeast respectively. The nearest HI-rich galaxy within 200 km s^{-1} is NGC 4192A at a projected distance of 290 kpc (Davies et al. 2004). If our detection were tidal debris, then it would have to have been drawn out on a timescale of $16 \text{ kpc}/200 \text{ km s}^{-1}$ (200 km s^{-1} being the typical velocity width within a single Arecibo beam), or $6 \times 10^7 \text{ yr}$. It follows that the interacting galaxies which generated it must still be close enough that they could have been near VIRGOHI21 $6 \times 10^7 \text{ yr}$ ago. For the two apparently nearest galaxies (above), that would imply relative projected speeds of greater than 1500 km s^{-1} . This is very high compared to the velocities of galaxies in the outskirts of Virgo (velocity dispersion $\sim 700 \text{ km s}^{-1}$) where VIRGOHI21 appears to be situated (Davies et al. 2004), and far too high to favor significant tidal interactions (e.g. Toomre & Toomre 1972; Barnes & Hernquist 1992). Furthermore, the observations of VIRGOHI21 show higher velocities to the north and lower velocities to the south, while the two nearby galaxies are placed with the one at a higher velocity to the south and the one with a lower velocity slightly north of due east – the opposite sense to that expected if the velocity gradient in VIRGOHI21 were due to a tidal interaction between them.

If a tidal origin is thus excluded what alternative explanations are compatible with the wide velocity width? Several narrow-line higher column-density clouds at different velocities lined up in the beam? Clouds like this are often associated with tidal debris as the filamentary structure breaks up into separate HI clouds which may form tidal dwarf galaxies (e.g. Hunsberger, Charlton & Zaritsky 1996). Such clouds should have been detected by our VLA observations; that they were not implies that this is an unlikely explanation. Another

possibility is that the gas is not bound – but then it should have dispersed in the same short time-scale of 6×10^7 yr. Given the dynamical timescale of the cluster of $\sim 10^9$ yr this possibility seems unlikely. The Galactic extinction at this point in the sky is only 0.15 mags in B band (Schlegel et al. 1998), therefore it is very unlikely that a galaxy has been hidden by obscuration. As the other possibilities seem so unlikely, one is left with the hypothesis that the gas in VIRGOHI21 is gravitationally bound and moving in stable, bound orbits which prevent shocking – rotation of course comes to mind, as in a flattened disk – a model which is not inconsistent with the spectra. If the system is bound, then its dynamical mass $M_{dyn} \equiv R_{HI} \times \Delta V^2 / G$ is greater than 9×10^{10} solar masses (with $R_{HI} \geq 8$ kpc and $\Delta V = 220$ km s $^{-1}$), not atypical of a rotating galaxy, though its $M_{dyn}/M_{HI} > 400$ is about 5 times higher than normal spirals.

From the well known Tully-Fisher correlation between rotational velocity and luminosity, calibrated in the Virgo Cluster by Fouqué et al. (1990), our HI detection, if indeed it is a bound system, should correspond to a galaxy with an absolute B magnitude of -19. At the distance of the Virgo Cluster this would correspond to a 12th magnitude galaxy, which would normally be extremely prominent at optical wavelengths. VIRGOHI21 appears to be a massive object not containing the expected bright galaxy.

It has been proposed that there is an HI column-density threshold ($\sim 10^{20}$ cm $^{-2}$) below which star formation ceases to occur (Toomre 1964; Martin & Kennicutt 2001). The mean column-density across our central beam at Arecibo is somewhat lower than this, at 4×10^{19} cm $^{-2}$ and our VLA observations set an upper limit to the column-density of 10^{20} cm $^{-2}$. This low column-density provides an explanation for the lack of an optical counterpart: this may be a dark galaxy that has failed to form stars because the low disk surface-density prevents fragmentation of the gas, i.e. it does not satisfy Toomre’s criterion (Verde et al. 2002; Toomre 1964).

If such dark objects exist in significant numbers, then why has it taken until now to detect one? VIRGOHI21-like objects could only have been detected by HI surveys which meet three criteria: (1) that they are ‘blind’, rather than targeted at previously-identified objects (which, by definition, are not ‘dark’); (2) that they have 5σ column-density sensitivity to galaxies with velocity widths ~ 200 km s $^{-1}$ at the 5×10^{19} cm $^{-2}$ level (which rules out older HI surveys); (3) that they have complete optical follow-up observations to deep isophotal limits. While there have been many blind surveys, the only ones to meet the second criterion are HIPASS (Meyer et al. 2004), HIJASS (Lang et al. 2003), AHISS (Zwaan et al. 1997), HIDEEP (Minchin et al. 2003) and VIRGOHI (Davies et al. 2004) and of these, only the last three satisfy the third criterion – that they have complete optical follow-up data. The HIPASS Bright Galaxy Catalogue (Koribalski et al. 2004) (BGC; peak flux > 116 mJy),

which has recently been used to make the most accurate determination to date of the HI mass function (Zwaan et al. 2003), would not have detected VIRGOHI21 unless it were within six Mpc – a relatively small distance and very close to the level where it would be impossible, in velocity-space, to distinguish a truly isolated HI cloud from one associated with The Galaxy. Taking the volume in which VIRGOHI21 would have been detected (in HIDEEP, VIRGOHI and AHISS) leads to a global density of $\sim 0.02 \text{ Mpc}^{-3}$, equivalent to a contribution to the cosmic density of $\Omega \simeq 0.01$. To have had more than one detection would therefore imply a very significant contribution to the cosmic density.

4. Conclusions

In the very nature of things it would be difficult to make an indisputable claim to have found a dark galaxy, particularly when past claims to that effect have quickly been ruled out by subsequent observations (either of a dim underlying galaxy or of bridging connections to nearby visible companions). Nevertheless, VIRGOHI21 passes all of the careful tests we have been able to set for it, using the best equipment currently available. Far longer VLA observations might help – but the very low column density and broad velocity width will make VIRGOHI21 an extremely challenging test for any current interferometer. And if every deep 21-cm detection without an optical counterpart is dismissed out of hand as debris – without considering the timing argument we present in Section 3 (which can be used to exclude all previous claims) – one is in effect ruling out, by definition, the detection of any dark galaxy at 21-cm. We cannot of course be certain, but VIRGOHI21 has turned up in one of the two extremely deep 21-cm surveys where you could most reasonably expect to find a dark galaxy, and meets all of the criteria we can, in practice, set for such an elusive but potentially vital object today: in particular such a broad observed velocity width cannot itself be attributed to a tidal interaction if the putative interacting galaxies are well outside the Arecibo beam – as they surely must be here. Future deep HI surveys could reveal a population of such galaxies; with colleagues we are planning such at Arecibo, Jodrell Bank and Parkes.

REFERENCES

- Barnes, J. E. & Hernquist, L. 1992, ARA&A, 30, 705
Benson, A. J., Frenk, C. S., Baugh, C. M., Cole, S. & Lacey, C. G. 2001, MNRAS, 327, 1041
Bothun, G. D., Impey, C. D., Malin, D. F. & Mould, J. R. 1987, AJ, 94, 23

- Boyce, P. J. et al. 2001, ApJ, 560, L127
- Davies, J. et al. 2004, MNRAS, 349, 922
- Fouqué, P., Bottinelli, L., Gouguenheim, L. & Paturel, G. 1990, ApJ, 349, 1
- Freeman, K. C. 1970, ApJ, 160, 811
- Giovanelli, R. & Haynes M. P. 1989, ApJ, 346, 5
- Graham, J. A. et al. 1999, ApJ, 516, 626
- Hunsberger, S. D., Charlton, J. C., & Zaritsky, D. 1996, ApJ, 462, 50
- Jimenez, R., Heavens, A. F., Hawkins, M. R. S. & Padoan, P. 1997, MNRAS, 292, L5
- Karachentsev, I. D., Karachentseva, V. E., Huchtmeier, W. K. & Makarov, D. I. 2004, AJ, 127, 2031
- Kilborn, V. A. et al. 2000, ApJ, 120, 1342
- Klypin, A., Kravtsov, A. V., Valenzuela, O. & Prada, F. 1999, ApJ, 522, 82
- Koribalski, B. S. et al. 2004, AJ, 128, 16
- Lang, R. et al. 2003, MNRAS, 342, 738
- McMahon, R. G., Irwin, M. J., Giovanelli, R., Haynes, M. P., Wolfe, A. M. & Hazard, C.. 1990, ApJ, 359, 302
- Martin C. L. & Kennicutt, R. C. 2001, ApJ, 555, 301
- Meyer, M. J. et al. 2004, MNRAS, 350, 1195
- Minchin, R. F. et al. 2003, MNRAS, 346, 787
- Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J. & Tozzi, P. 1999, ApJ, 524, L19
- Roberts, S. et al. 2004, MNRAS, 352, 478
- Ryder, S. D. et al. 2001, ApJ, 555, 232
- Sabatini, S., Davies, J., Scaramella, R., Smith, R., Baes, M., Linder, S. M., Roberts, S., & Testa, V. 2003, MNRAS, 341, 981

- Salucci, P. & Persic, M. 1997, in ASP Conf. Ser. 117, Dark and Visible Matter in Galaxies, ed. M. Persic & P. Salucci (San Francisco: ASP), 242
- Schlegel, D. J., Finkbeiner, D. P. & Davis, M. 1998, ApJ, 500, 525
- Schneider, S. E., Helou, G., Salpeter, E. E. & Terzian, Y., 1983, ApJ, 273, 1
- Spergel, D. N. et al. 2003, ApJ, 148, 175
- Toomre, A. & Toomre, J. 1972, ApJ, 178, 623
- Toomre, A. 1964, ApJ, 139, 1217
- Verde, L., Oh, S. P. & Jimenez, R. 2002, MNRAS, 336, 541
- Zucker, D. B. et al. 2004, ApJ, 612, L121
- Zwaan, M. A., Briggs, F. H., Sprayberry, D. & Sorar, E. 1997, ApJ, 490, 173
- Zwaan, M. A. et al. 2003, AJ, 125, 2842




Fig. 1.— Atomic Hydrogen spectra of VIRGOHI21 plotted as a function of recessional velocity. (a) discovery spectrum from Jodrell Bank (3500 s). (b) - (d) Arcibo follow-up spectra from beams labelled (b) - (d) in Fig. 2, cutting north-south across the source (600 s). (e) Sum of all Arcibo spectra (600 s per beam). The peak at $\sim 1750 \text{ km s}^{-1}$ in spectrum (e) can be clearly associated with galaxy C on Fig. 3 by our VLA data.




Fig. 2.— The Arecibo pointing pattern for VIRGOHI21 overlaid on a digitized sky survey image of the region around the source. Circles mark the beam position and FWHM; black circles denote beams where a firm detection (better than 4σ) was made and grey circles those beams which do not contain a definite signal. The large circle marks the beam corresponding to the pixel in the Jodrell Bank data cube from which the spectrum and measurements in Fig. 1 (a) are taken. The beams marked (b), (c) and (d) correspond to the spectra marked (b), (c) and (d) in Fig. 1




Fig. 3.— Isaac Newton Telescope *B*-band optical image (shown as negative) of the field of VIRGOHI21. The cross marks the weighted center of the HI detection and the circle shows the size and position of the central Arcibo beam. Optical sources labelled (A) – (E) are discussed in the text. Contours show the moment map from the VLA detection of source (C) ($M_{HI} = 1.4 \times 10^7 M_{\odot}$) at 1750 km s^{-1} and are at 3, 6, 9 and 12 sigma.