

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/278660128>

Back to the future: the sterile insect technique against mosquito disease vectors

Article in *Current Opinion in Insect Science* · August 2015

DOI: 10.1016/j.cois.2015.05.011

CITATIONS

119

READS

2,345

5 authors, including:



Rosemary Lees

Liverpool School of Tropical Medicine

12 PUBLICATIONS 265 CITATIONS

[SEE PROFILE](#)



Jeremie R L Gilles

Insects For Better Future

135 PUBLICATIONS 2,508 CITATIONS

[SEE PROFILE](#)



Jorge Hendrichs

International Atomic Energy Agency (IAEA)

104 PUBLICATIONS 4,902 CITATIONS

[SEE PROFILE](#)



Marc J B Vreysen

International Atomic Energy Agency (IAEA)

147 PUBLICATIONS 3,609 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Potential of the SIT for controlling *Drosophila suzukii* [View project](#)



male sex determination, transcriptomics, embryos, functional genomics [View project](#)

Back to the future: the sterile insect technique against mosquito disease vectors

Rosemary Susan Lees, Jeremie RL Gilles, Jorge Hendrichs, Marc JB Vreysen and Kostas Bourtzis



With the global burden of mosquito-borne diseases increasing, and some conventional vector control tools losing effectiveness, the sterile insect technique (SIT) is a potential new tool in the arsenal. Equipment and protocols have been developed and validated for efficient mass-rearing, irradiation and release of *Aedines* and *Anophelines* that could be useful for several control approaches. Assessment of male quality is becoming more sophisticated, and several groups are well advanced in pilot site selection and population surveillance. It will not be long before SIT feasibility has been evaluated in various settings. Until perfect sexing mechanisms exist, combination of *Wolbachia*-induced phenotypes, such as cytoplasmic incompatibility and pathogen interference, and irradiation may prove to be the safest solution for population suppression.

Address

Insect Pest Control Sub-programme, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Vienna, Austria

Corresponding author: Bourtzis, Kostas (K.Bourtzis@iaea.org)

Current Opinion in Insect Science 2015, 10:156–162

This review comes from a themed issue on **Vectors and medical and veterinary entomology**

Edited by **Nora Besansky**

<http://dx.doi.org/10.1016/j.cois.2015.05.011>

2214-5745/© 2015 The Authors. Published by Elsevier Inc.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The many pathogens transmitted by mosquitoes (Diptera: Culicidae) which feed on the blood of humans in order to mature their eggs are responsible for enormous suffering worldwide. Annual deaths from malaria alone number at least 600,000, up to 100,000 people contract dengue each year, and Chikungunya causes severe chronic joint pain in patients across the globe (World Health Organization factsheet; URL: <http://www.who.int/mediacentre/factsheets/fs387/en/>). Aside from causing mortality and morbidity, the economic and social burden from these diseases is significant [65], particularly in SubSaharan Africa (Multisectoral Action Framework for Malaria; URL: <http://reliefweb.int/sites/>

reliefweb.int/files/resources/Multisectoral-Action-Framework-for-Malaria.pdf).


The pressure placed on humanity by these vectors is increasing. Expansion of the distribution of several invasive *Aedes* species such as *Aedes albopictus* [1] is evident in many areas, including Europe [2,3] and USA [4]. Modelling and field experiments have predicted that *Ae. albopictus* has the potential to invade large areas of Australia [5] and urbanisation is increasing its abundance in China [6]. With no effective vaccines or specific drugs to prevent or treat mosquito-borne diseases, the best line of defence is to combat the vector, to remove the contact between mosquitoes and humans and thus interrupt the disease transmission cycle. Effective mosquito control is hindered by growing insecticide resistance of malaria [7] and dengue vectors [8], even in regions only recently invaded (e.g. [9]). There is therefore increasing demand for complementary tactics that are effective, more sustainable and friendly to the environment.

One such tactic could be the sterile insect technique (SIT), which relies on the production and release of sufficient sterile males to induce sterility in the wild females which, over time, causes the target population to decline (Figure 1). The SIT has no regulatory requirements and the technique would be combined with others as part of an area-wide integrated pest management (AW-IPM) approach to reduce the vector population below the threshold required for disease transmission. Sterilisation using ionising radiation has been extremely effective and applied successfully for population suppression, containment or eradication of several major pest insect species [10].

Rather than sterilising males using irradiation, an alternative method is to exploit the natural phenomenon of cytoplasmic incompatibility (CI). In most diplo-diploid species, CI is expressed as embryonic lethality after matings between *Wolbachia*-infected males and uninfected females or females infected with a different *Wolbachia* strain [11]. Proof-of-concept has been provided that CI could be used to manage agricultural pests and disease vectors through population suppression or replacement approaches [12,13,14]. CI-based population suppression is known as the incompatible insect technique (IIT) (Figure 1).

As the key mosquito disease vectors are all relatively amenable to colonisation and rearing, and in many

Figure 1



The sterile insect technique (SIT), incompatible insect technique (IIT) or a combination of the two could be used to suppress mosquito populations. Male mosquitoes are sterilised either by the application of irradiation or (trans)infection with *Wolbachia*, or both, and then released into the target population to sterilise the wild females.

situations the natural population densities are low, the SIT, IIT, or a combination of the two are well suited for their management. The advantages of combining these tactics will be discussed in this review, alongside the current state of the art for the two approaches. Much progress has been made in recent years towards developing the required technology and methodology to bring mosquito suppression using sterility to field application; indeed pilot releases have begun in a number of sites around the world. It should be mentioned, however, that a number of other technologies have also been developed and are being tested in pilot trials including RIDL (Release of Insects carrying a Dominant Lethal) and *Wolbachia*-based population replacement strategies. However, it is beyond the scope of this review to discuss these approaches, and they have recently been reviewed elsewhere [13,15].

Developing the sterile insect technique against mosquitoes

After a period of enthusiasm in the 1960s to early 1980s [16], the use of sterile male release for mosquito control was largely abandoned. However, the growing pressures

from mosquito-borne pathogens described above, and the proposed use of modern biotechnologies to sterilise or otherwise alter mosquitoes, have led to revived interest in recent years.

In the last decade, the Joint FAO/IAEA Programme and their collaborators have been the main drivers for the development of the “SIT package” for mosquitoes. Requests from many countries to develop and evaluate the SIT for use against mosquitoes have spurred the development and ongoing validation of mass-rearing equipment, diet and protocols for *Anopheles* and *Aedes* species. Diets have been optimised to feed the larval stages of *An. gambiae* [17], *An. arabiensis* [18], *An. stephensi* [19] and *Ae. albopictus* [20]. *Anopheles* larvae can be mass-reared efficiently in large trays fitted into a novel tilting rack system [21,22], and the system is being validated for *Aedes* species. Anopheline pupae can be separated from larvae based on differential buoyancy using custom vortex equipment [23], or the Fay-Morlan separator used for *Aedes* [24], quantified volumetrically, and allowed to emerge into adult mass-rearing cages [25]. Blood meals are offered to females using a modified hemotec membrane feeder and

water is added to the cage floor for oviposition [26]. Eggs are flushed from these cages and can be quantified ([27] for *Aedes*, Maiga *et al.*, personal communication for *Anopheles*), stored and hatched effectively [28] to give a reliable quantity of eggs and so a predictable larval density in rearing trays.

A method to accurately separate males from females on a large scale, crucial for the required male-only release, is still required [29], particularly for *Anopheles*, which do not have the sexual dimorphism that allows the sexing of *Aedes* on a small-medium scale [24,30^{*}]. A method to spike blood meals with Ivermectin [31] is a reasonable stop-gap solution. This requirement for sex separation applies for all mosquito suppression methods based on release, including those described below. The methods for radiation-sterilisation of mosquitoes have long been available, developed alongside those for many other target species, but have more recently been revisited to optimise doses [32–34], and to assess the use of X-rays as an alternative for gamma irradiation [35,36].

Progress in SIT field application

The vanguard in reviving the use of the SIT against mosquitoes was an Italian group [37] who released around 1000 irradiated *Ae. albopictus* pupae per hectare per week, inducing up to 68% sterility in the target populations in three pilot sites of between 16 and 45 ha [38^{*}]. Releases continued for 5 years, and demonstrated the potential of sterile males to suppress an *Ae. albopictus* population.

The importance of quality management of sterile mosquito males to ensure adequate performance and competitiveness after release is evident from examples in other species [39]; the estimation and quantification of the impact of mass-rearing, radiation and handling on male mating competitiveness of sterile males has attracted a lot of research. Semi-field and field experiments have demonstrated that a radiation dose can be selected that gives sufficient sterility without significantly impacting competitiveness [40,41,42^{*}]. With this reassurance, several vector control groups, supported by the FAO/IAEA, are conducting preparatory activities and initiating pilot trials that include the SIT.

The first step in assessing the SIT in a given context is to select proper pilot sites, which should have a manageable size, a low mosquito population density and a good level of geographical or biological isolation, among other criteria reviewed in Malcolm *et al.* [43] and Brown *et al.* [44]. Two such sites have been identified by the Ministry of Health and Quality of Life in Mauritius [45], where they have also completed the second preparatory step, the long-term surveillance of the *Aedes albopictus* natural population using ovitraps and BG-Sentinels to trap adults. A project in La Réunion has progressed to a similar stage [46]. A good understanding of the biology, dynamics and

distribution of the male population in the target area is crucial to properly plan the releases and to monitor their effect. Although several effective traps exist for male *Aedes* surveillance, male *Anopheles* are much more difficult to monitor.

In Sudan, the target species, *An. arabiensis*, is contained along a narrow strip on either side of the River Nile, and surveillance has demonstrated temporal variations of population densities that were overall low [47]. Further, mark-release-recapture experiments have demonstrated the ability of radiation-sterilised males to locate and participate in naturally occurring swarms, or to start new swarms [42^{*}]. Encouraged by these data, small-scale releases have started, and construction of a mass-rearing facility is scheduled to supply the sterile mosquitoes for suppression trials. A project in South Africa, targeting *An. arabiensis*, is at a similar stage of advancement [48]. A coordinated research project (CRP) is being initiated by the FAO/IAEA (“Mosquito Handling, Transport, Release and Male Trapping Methods”) to support these projects in developing and validating suitable methods for releasing sterile male mosquitoes and surveying the target population before, during and after suppression trials (<http://www-naweb.iaea.org/nafa/ipc/index.html>).

With these pieces in place, the whole SIT package for mosquitoes is coming closer to full scale field trials, and it is expected that within a very few years multiple feasibility studies in a range of settings and against a number of species will have been completed and have demonstrated the effectiveness and applicability of the technique against these disease vectors.

Incompatible insect technique: an additional tool and its potential combination with the sterile insect technique for population suppression

About 50 years ago *Wolbachia*-induced CI as a tool to suppress natural populations of *Culex pipiens fatigans* was used for the first time [49]. During the last few years there have been significant developments, both in the laboratory and in the field, towards the use of IIT for population suppression of mosquito vectors. There are also self-sustaining *Wolbachia*-based approaches that target population replacement with CI-inducing and pathogen-blocking strains; however the applicability, effectiveness and sustainability of this strategy require more studies [13,50].

Wolbachia-infected lines of *Culex pipiens quinquefasciatus* were selected and tested in laboratory cages for CI expression and population suppression of four natural populations originating from four islands: Grand Glorieuse, Mauritius, Mayotte and La Réunion [51^{*},52^{*}]. The results of these trials were very promising, indicating that

C. p. quinquefasciatus males infected with the *Wolbachia* strain *wPip*(Is) were fully incompatible (100% CI) with females from the four islands of the south-western Indian Ocean [51^{*}]. As a next step, semi-field experiments were run showing that the *wPip*(Is) males were: (a) able to induce complete CI in La Réunion field females and (b) fully competitive against field-collected males to mate with field-collected females [52^{*}]. Similar IIT-based pilot trails were implemented against *Aedes polynesiensis*, with a proof-of-concept pilot trial in the Society Islands [53,54], and an ongoing field trial in French Polynesia (Bossin, personal communication).

New *Wolbachia* infection and CI types have been developed for one of the major dengue vectors, *Aedes albopictus* [55]. One of the lines, ARwP, is infected with a *wPip* strain which naturally occurs in *Culex pipiens*. Mating experiments have shown that ARwP males exhibited full CI with uninfected females or naturally double-infected females (*wAlbA* and *wAlbB*), suggesting that this strain could in principle be used for population suppression. The use of ARwP (*wPip*) males to suppress naturally double-infected (*wAlbA* and *wAlbB*) *Ae. albopictus* populations would be advantageous if there was complete CI between these strains, to minimise the risk from any accidental release of *wPip*-infected females. However a recent study showed that crosses between males with low *wAlbA* density and ARwP females were partially fertile [56]. This finding suggests that the accidental release of *wPip*-infected ARwP females may jeopardise a population suppression programme by instead causing population replacement. Thus IIT application will require a fail-proof sexing method so that it could be used as a tactic to suppress populations of mosquito vector species in a way similar to the SIT.

The requirement for perfect separation of males and females prior to release, discussed above, is particularly important for IIT because the accidental release of females may result in the loss of IIT and render a population suppression programme into population replacement. A possible strategy to manage this risk is to combine SIT and IIT (Figure 1) [14,53,57,58]. A strategy combining a low radiation dose to ensure female sterility (SIT), and IIT is being initiated against *Ae. albopictus*. A triple-infected line (*wAlbA*, *wAlbB* and *wPip*) was shown to be completely incompatible with double-infected (*wAlbA* and *wAlbB*) lines as well as providing protection against dengue (Xi, unpublished data). No significant negative impact of the triple infection on several fitness traits could be measured [59], though the time required for immature development was significantly reduced in males compared to females, a finding which could be explored for sex separation.

Until recently, *Anopheles* species were considered to be *Wolbachia*-free. However, *Wolbachia* was recently

detected in a natural population of *Anopheles gambiae* in Burkina Faso [60]. The *wAlbB* strain was recently transferred from *Ae. albopictus* to *Anopheles stephensi* creating a new stable transinfected line expressing complete CI, produced for population replacement strategies, but potentially effective for population suppression [13]. Taken together, these data suggests that *Anopheles* species are not “resistant” to *Wolbachia* infection and that IIT could also be used for population suppression of Anophelines.

Conclusion

In response to the growing interest and demand for the development and application of the SIT, with possible combination with the related IIT against mosquito vectors, significant advances have been made in developing the required equipment and protocols for rearing, sterilising and assessing the quality of male *Aedes* and *Anopheles* mosquitoes. Most of the pieces are thus in place for the technique to be validated in suppression programmes on a small scale, and in several different settings the preliminary work of site selection, population surveillance, up-scaling of rearing and quality control is well advanced.

Before large scale releases are feasible, however, more efficient and less labour intensive methods are needed for transporting and releasing male mosquitoes into the field, as well as more effective methods for monitoring programme progress, particularly for *Anopheles* species. These are fairly simple design and engineering questions, which some time and careful evaluation in the field will be able to address in due course. The other major challenge is the development of an accurate sex separation method which can be applied on a large scale, which may require more sophisticated developments [29]. Until perfect sexing is available, the combination of SIT and IIT could be used to ensure that any unintentionally released *Wolbachia*-infected females would be sterile. In addition, *Wolbachia* transinfection may provide protection against the establishment, replication and/or transmission of *Plasmodium*, also eliminating the risk of disease transmission [13,14,61]. Once these remaining pieces are in place, sterile male release programmes hold great promise for control of mosquito vectors, particularly in urban areas where the human population to be protected is concentrated.

The effects of any genetic manipulation on the robustness and competitiveness of male mosquitoes in an open field setting is difficult to know before release. It is likely that genetic modification will have an impact [62], though the extent of the effect will be strain-specific. *Wolbachia* transinfection may or may not negatively impact mosquitoes [59,63], and may not interrupt disease transmission. With the application of radiation it is possible to adapt the dose to induce an adequate level of sterility whilst minimising the effect on male performance. The random mutations and gross gonad damage caused by irradiation

[16] eliminate the risk of resistance, which is a major problem with insecticide use, and potentially with genetic control measures [64]. Finally, in circumstances where there is public or regulatory opposition to the use of genetically modified organisms, release of fertile *Wolbachia*-infected females for population replacement, or generic insecticides, the SIT and the SIT–IIT combination offer an acceptable alternative. It is hoped that the effectiveness of these techniques can be demonstrated in the near future, and if proved effective, another powerful tool will have been added to the limited arsenal available for use against mosquito-borne diseases.

Acknowledgements

The authors apologise for not citing many colleagues' work due to limited space. The authors would also like to thank colleagues for useful discussions in the framework of Joint FAO/IAEA meetings.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
1. Bonizzoni M, Gasperi G, Chen X, James AA: **The invasive mosquito species *Aedes albopictus*: current knowledge and future perspectives.** *Trends Parasitol* 2013, **29**:460-468.
 2. Schaffner F, Mathis A: **Dengue and dengue vectors in the WHO European region: past, present, and scenarios for the future.** *Lancet Infect Dis* 2014, **14**:1271-1280.
 3. Medlock JM, Hansford KM, Versteirt V, Cull B, Kampen H, Fontenille D, Hendrickx G, Zeller H, Van Bortel W, Schaffner F: **An entomological review of invasive mosquitoes in Europe.** *Bull Entomol Res* 2015, **25**:1-27.
 4. Rochlin I, Ninivaggi DV, Hutchinson ML, Farajollahi A: **Climate change and range expansion of the Asian tiger mosquito (*Aedes albopictus*) in Northeastern USA: implications for public health practitioners.** *PLOS ONE* 2013, **8**:e60874.
 5. Nicholson J, Ritchie SA, Russell RC, Zalucki MP, Van Den Hurk AF: **Ability for *Aedes albopictus* (Diptera: Culicidae) to survive at the climatic limits of its potential range in eastern Australia.** *J Med Entomol* 2014, **51**:948-957.
 6. Li Y, Kamara F, Zhou G, Puthiyakunnon S, Li C, Liu Y, Zhou Y, Yao L, Yan G, Chen X-G: **Urbanization increases *Aedes albopictus* larval habitats and accelerates mosquito development and survivorship.** *PLoS Negl Trop Dis* 2014, **8**:e3301.
 7. Sokhna C, Ndiath MO, Rogier C: **The changes in mosquito vector behaviour and the emerging resistance to insecticides will challenge the decline of malaria.** *Clin Microbiol Infect* 2013, **19**:902-907.
 8. Ranson H, Burhani J, Lumjuan N, Black WC IV: **Insecticide resistance in dengue vectors.** *TropIKAet* 2010, **1**:1-12.
 9. Marcombe S, Farajollahi A, Healy SP, Clark GG, Fonseca DM: **Insecticide resistance status of United States populations of *Aedes albopictus* and mechanisms involved.** *PLOS ONE* 2014, **9**:e101992.
 10. Dyck VA, Hendrichs J, Robinson AS: *Sterile insect technique: principles and practice in area-wide integrated pest management.* Dordrecht, The Netherlands: Springer; 2005.
 11. Saridaki A, Bourtzis K: ***Wolbachia*: more than just a bug in insects genitals.** *Curr Opin Micro* 2010, **13**:67-72.
 12. Bian G, Joshi D, Dong Y, Lu P, Zhou G, Pan X, Xu Y, Dimopoulos G, Xi Z: ***Wolbachia* invades *Anopheles stephensi* populations and induces refractoriness to *Plasmodium* infection.** *Science* 2013, **340**:748-751.
- This study presents the successful transfection of the malaria vector *Anopheles stephensi* with the *Wolbachia* strain wAlbB, the perfect maternal transmission of the symbiont, the induction of high levels of cytoplasmic incompatibility and associated invasion dynamics, and the symbiont-mediated resistance against the malaria parasite *Plasmodium falciparum*.
13. McGraw EA, O'Neill SL: **Beyond insecticides: new thinking on an ancient problem.** *Nat Rev Microbiol* 2013, **11**:181-193.
 14. Bourtzis K, Dobson SL, Xi Z, Rasgon JL, Calvitti M, Moreira LA, Bossin HC, Moretti R, Baton LA, Hughes GL, Mavingui P, Gilles JRL: **Harnessing mosquito-*Wolbachia* symbiosis for vector and disease control.** *Acta Trop* 2014, **132**:S150-S163.
 15. Alphey L: **Genetic control of mosquitoes.** *Ann Rev Entomol* 2014, **59**:205-224.
 16. Klassen W, Curtis CF: **History of the sterile insect technique.** In *Sterile insect technique: principles and practices in area-wide integrated pest management.* Edited by Dyck VA, Hendrichs J, Robinson AS. Springer; 2005:39-68.
 17. Yahouedo GA, Djogbenou L, Saizonou J, Assogba BS, Makoutode M, Gilles JRL, Maiga H, Mouline K, Soukou BK, Simard F: **Effect of three larval diets on larval development and male sexual performance of *Anopheles gambiae* s.s.** *Act Trop* 2014, **132**:S96-S101.
 18. Damiens D, Benedict MQ, Wille M, Gilles JRL: **An inexpensive and effective larval diet for *Anopheles arabiensis* (Diptera: Culicidae): eat like a horse, a bird or a fish?** *J Med Entomol* 2012, **49**:1001-1011.
 19. Khan I, Farid A, Zeb A: **Development of inexpensive and globally available larval diet for rearing *Anopheles stephensi* (Diptera: Culicidae) mosquitoes.** *Parasit Vectors* 2013, **6**:90.
 20. Puggioli A, Balestrino F, Damiens D, Lees RS, Soliban SM, Madakacherry O, Dindo ML, Bellini R, Gilles JRL: **Efficiency of three diets for larval development in mass rearing *Aedes albopictus* (Diptera: Culicidae).** *J Med Entomol* 2013, **50**:819-825.
 21. Balestrino F, Benedict MQ, Gilles JR: **A new larval tray and rack system for improved mosquito mass rearing.** *J Med Entomol* 2012, **49**:595-605.
 22. Balestrino F, Puggioli A, Gilles JRL, Bellini R: **Validation of a new larval rearing unit for *Aedes albopictus* (Diptera: Culicidae) mass rearing.** *PLOS ONE* 2014, **9**:e91914.
 23. Balestrino F, Giles JRL, Soliban SM, Nirschl A, Benedict QE, Benedict MQ: **Mosquito mass rearing technology: a cold-Water vortex device for continuous unattended separation of *Anopheles arabiensis* pupae from larvae.** *J Am Mosq Cont Assoc* 2011, **27**:227-235.
 24. Focks DA: **An improved separator for the developmental stages, sexes, and species of mosquitoes (Diptera: Culicidae).** *J Med Entomol* 1980, **17**:567-568.
 25. Balestrino F, Puggioli A, Bellini R, Petric D, Gilles JRL: **Mass production cage for *Aedes albopictus* (Diptera: Culicidae).** *J Med Entomol* 2014, **51**:155-163.
 26. Damiens D, Soliban SM, Balestrino F, Alsir R, Vreysen MJB, Gilles JRL: **Different blood and sugar feeding regimes affect the productivity of *Anopheles arabiensis* colonies (Diptera: Culicidae).** *J Med Entomol* 2013, **50**(2):336-343 <http://dx.doi.org/10.1603/ME12212>.
 27. Zheng ML, Zhang DJ, Damiens DD, Yamada H, Gilles JRL: **Standard operating procedures for standardized mass rearing of the dengue and chikungunya vectors *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) – I – egg quantification.** *Parasit Vectors* 2015, **8**:42.
 28. Zheng ML, Zhang DJ, Damiens DD, Lees RS, Gilles JRL: **Standard operating procedures for standardized mass rearing of the dengue and chikungunya vectors *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) – II – egg storage and hatching.** *Parasit Vectors* 2015. [accepted].
 29. Gilles JRL, Schetelig MF, Scolari F, Marec F, Capurro ML, Franz G, Bourtzis K: **Towards mosquito sterile insect technique**

programmes: exploring genetic, molecular, mechanical and behavioural methods of sex separation in mosquitoes. *Act Trop* 2014, **132S**:S178-S187.

30. Carvalho DO, Nimmo D, Naish N, McKemey AR, Gray P, Wilke AB, Marrelli MT, Virginio JF, Alphey L, Capurro ML: **Mass production of genetically modified *Aedes aegypti* for field releases in Brazil.** *J Vis Exp* 2014, **83**:e3579.

A guide to the production of genetically modified *Aedes aegypti* is presented, including considerations in designing a suitable insectary and quality control of the produced male mosquitoes. The multimedia nature of this publication, which includes a video element, makes it a comprehensive and easy to use set of practical guidelines for the large scale rearing of *Aedes aegypti*.

31. Yamada H, Soliban SM, Vreysen MJB, Chadee DD, Gilles JRL: **Eliminating female *Anopheles arabiensis* by spiking blood meals with toxicants as a sex separation method in the context of the sterile insect technique.** *Parasit Vect* 2013, **6**:197.
32. Helinski MEH, Parker AG, Knols BGJ: **Radiation-induced sterility for pupal and adult stages of the malaria mosquito *Anopheles arabiensis*.** *Malar J* 2006, **5**:41.
33. Balestrino F, Medici A, Candini G, Carrieri M, Maccagnani B, Calvitti M, Maini S, Bellini R: **Gamma ray dosimetry and mating capacity studies in the laboratory on *Aedes albopictus* males.** *J Med Entomol* 2010, **47**:581-591.
34. Yadav K, Dhiman S, Baruah I, Singh L: **Effect of gamma radiation on survival and fertility of male *Anopheles stephensi* liston, irradiated as pharate adults.** *J Ecobiotech* 2010, **2**:6-10.
35. Ndo C, Yamada H, Damiens DD, N'do A, Seballos G, Gilles JRL: **X-ray sterilization of the *An. arabiensis* genetic sexing strain 'ANOIPCL1' at pupal and adult stages.** *Acta Trop* 2014, **131**:124-128.
36. Yamada H, Parker AG, Oliva CF, Balestrino F, Gilles JRL: **X-ray-induced sterility in *Aedes albopictus* (Diptera: Culicidae) and male longevity following irradiation.** *J Med Entomol* 2014, **51**:811-816.
37. Bellini R, Calvitti M, Medici A, Carrieri M, Celli G, Maini S: **Use of the sterile insect technique against *Aedes albopictus* in Italy: first results of a pilot trial.** In *In area-wide control of insect pests: from research to field implementation*. Edited by Vreysen MJB, Robinson AS, Hendrichs J. 2007:505-515.
38. Bellini R, Medici A, Puggioli A, Balestrino F, Carrieri M: **Pilot field trials with *Aedes albopictus* irradiated sterile males in Italian urban areas.** *J Med Entomol* 2013, **50**:317-325.

This is the first demonstration of the sterile insect technique against mosquitoes in recent times, and the best documented. Data from 3 years of weekly releases of radiation-sterilised *Aedes albopictus* during the peak summer months. A total of 2 million males were released that induced up to 68% sterility in the native population. Results suggest that a reduction in egg fertility of 70–80% is required to achieve a similar reduction in egg density in ovitraps.

39. Calkins CO, Parker AG: **Sterile insect quality.** In *In sterile insect technique: principles and practices in area-wide integrated pest management*. Edited by Dyck VA, Hendrichs J, Robinson AS. Springer; 2005:269-296.
40. Bellini R, Balestrino F, Medici A, Gentile G, Veronesi R, Carrieri M: **Mating competitiveness of *Aedes albopictus* radio-sterilized males in large enclosures exposed to natural conditions.** *J Med Entomol* 2013, **50**:94-102.
41. Madakacherry O, Lees RS, Gilles JRL: ***Aedes albopictus* (Skuse) males in laboratory and semi-field cages: release ratios and mating competitiveness.** *Act Trop* 2014, **132S**:124S-129S.
42. Ageep TB, Damiens D, Alsharif B, Ahmed A, Salih EHO, Ahmed FTA, Diabaté A, Lees RS, Gilles JRL: **El Sayed BB: participation of irradiated *Anopheles arabiensis* males in swarms following field release in Sudan.** *Malar J* 2014, **13**:484.

This study demonstrated the participation of radiation-sterilised male *Anopheles arabiensis* in naturally occurring swarms in Northern Sudan at a distance of up to 200 m within 2 hours of release. Competition with wild males for mates is critical to the success of the sterile insect technique, and this was a key demonstration of their ability to participate in natural courtship behaviour in the field. Results also suggest that irradiated males

may be able to initiate swarms, and point to the inclusion of a pre-release period as a means to increase mating competitiveness.

43. Malcolm CA, El Sayed B, Babiker A, Girod R, Fontenille D, Knols BGJ, Nugad AH, Benedict MQ: **Field site selection: getting it right first time around.** *Malar J* 2009, **8**:S9.
44. Brown DM, Alphey LS, McKemey A, Beech C, James AA: **Criteria for identifying and evaluating candidate sites for open-field trials of genetically engineered mosquitoes.** *Vector Borne Zoonotic Dis* 2014, **14**:291-299.
45. Iyaloo DP, Elahee KB, Bheecarry A, Lees RS: **Guidelines to site selection for population surveillance and mosquito control trials: a case study from Mauritius.** *Act Trop* 2014, **132**:S140-S149.
46. Boyer S, Toty C, Jacquet M, Lempérière G, Fontenille D: **Evidence of multiple inseminations in the field in *Aedes albopictus*.** *PLoS ONE* 2012, **7**:e42040 <http://dx.doi.org/10.1371/journal.pone.0042040>.
47. Ageep TB, Cox J, Hassan MM, Knols BGJ, Benedict MQ, Malcolm CA, Babiker A, El Sayed BB: **Spatial and temporal distribution of the malaria mosquito *Anopheles arabiensis* in northern Sudan: influence of environmental factors and implications for vector control.** *Malar J* 2009, **9**:123.
48. Munchenga G, Brooke BD, Spillings B, Essop L, Hunt RH, Midzi S, Govender D, Braack L, Koekemoer LL: **Field study site selection, species abundance and monthly distribution of anopheline mosquitoes in the northern Kruger National Park, South Africa.** *Malaria J* 2014, **13**:1-11.
49. Laven H: **A possible model for speciation by cytoplasmic isolation in the *Culex pipiens* complex.** *Bull World Health Organ* 1967, **37**:263-266.
50. Hoffmann AA, Iturbe-Ormaetxe I, Callahan AG, Phillips BL, Billington K, Axford JK, Montgomery B, Turley AP, O'Neill SL: **Stability of the *wMel* *Wolbachia* infection following invasion into *Aedes aegypti* populations.** *PLoS Negl Trop Dis* 2014, **8**:e3115.
51. Atyame CM, Pasteur N, Dumas E, Tortosa P, Tantely ML, Pocquet N, Licciardi S, Bheecarry A, Zumbo B, Weill M *et al.*: **Cytoplasmic incompatibility as a means of controlling *Culex pipiens quinquefasciatus* mosquito in the islands of the south-western Indian Ocean.** *PLoS Negl Trop Dis* 2011, **5**:e1440.
- See below.
52. Atyame CM, Cattel J, Lebon C, Flores O, Dehecq JS, Weill M, Gouagna LC, Tortosa P: ***Wolbachia*-based population control strategy targeting *Culex quinquefasciatus* mosquitoes proves efficient under semi-field conditions.** *PLOS ONE* 2015, **10**:e2880119.
- The above two studies showed that incompatible insect technique, which is based on *Wolbachia*-induced CI, can be used as tool to control populations of *Culex quinquefasciatus*. They indicated that LR[wPip(Is)] males are incompatible with target field females from La Réunion (and other islands from the South-Western Indian Ocean). Competition with wild males for mates is critical to the success of the incompatible insect technique, as is for the sterile insect technique. The second study by Atyame and colleagues shows that the LR[wPip(Is)] males successfully compete with field males in mating with field females.
53. Brelsfoard CL, Séchan Y, Dobson SL: **Interspecific hybridization yields strategy for South Pacific Filariasis vector elimination.** *PLoS Negl Trop Dis* 2008, **2**:e129.
54. O'Connor L, Plichart C, Sang AC, Brelsfoard CL, Bossin HC, Dobson SL: **Open release of male mosquitoes infected with a *wolbachia* biopesticide: field performance and infection containment.** *PLoS Negl Trop Dis* 2012, **6**:e1797.
55. Calvitti M, Moretti R, Skidmore AR, Dobson SL: ***Wolbachia* strain wPip yields a pattern of cytoplasmic incompatibility enhancing a *Wolbachia*-based suppression strategy against the disease vector *Aedes albopictus*.** *Parasit Vectors* 2012, **5**:254.
56. Calvitti M, Marini F, Desiderio A, Puggioli A, Moretti R: ***Wolbachia* density and cytoplasmic incompatibility in *Aedes albopictus*: concerns with using artificial *Wolbachia* infection as a vector suppression tool.** *PLoS ONE* 2015, **10**:e0121813.

57. Arunachalam N, Curtis CF: **Integration of radiation with cytoplasmic incompatibility for genetic control in the *Culex pipiens* complex (Diptera: Culicidae).** *J Med Entomol* 1985, **22**:648-653.
58. Bourtzis K, Robinson AS: **Insect pest control using *Wolbachia* and/or radiation.** In *Insect symbiosis*, vol. 2. Edited by Bourtzis K, Miller T. CRC Press; 2006:225-246.
59. Zhang D, Zheng X, Xi Z, Bourtzis K, Gilles JRL: **Combining the sterile insect technique with the incompatible insect technique: I – Impact of *Wolbachia* infection on the fitness of triple- and double-infected strains of *Aedes albopictus*.** *PLOS ONE* 2015, **10**:e0121126.
60. Baldini F, Segata N, Pompon J, Marcenac P, Robert Shaw W, Dabiré RK, Diabaté A, Levashina EA, Catteruccia F: **Evidence of natural *Wolbachia* infections in field populations of *Anopheles gambiae*.** *Nat Commun* 2014, **5**:3985.
61. Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, Rocha BC, Hall-Mendelin S, Day A, Riegler M, Hugo LE *et al.*: **A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and Plasmodium.** *Cell* 2009, **139**:1268-1278.
62. Marrelli MT, Moreira CK, Kelly D, Alphey L, Jacobs-Lorena M: **Mosquito transgenesis: what is the fitness cost?** *Trends Parasitol* 2006, **22**:197-202.
63. Yeap HL, Mee P, Walker T, Weeks AR, O'Neill SLO, Johnson P, Ritchie SA, Richardson KM, Doig C, Endersby NM, Hoffmann AA: **Dynamics of the “popcorn” *Wolbachia* infection in outbred *Aedes aegypti* informs prospects for mosquito vector control.** *Genetics* 2011, **187**:583-595.
64. Alphey N, Bonsall MB, Alphey L: **Modeling resistance to genetic control of insects.** *J Theor Biol* 2011, **270**:42-55.
65. Shepard DS, Undurraga EA, Betancourt-Cravioto M, Guzmán MG, Halstead SB, Harris E, Mudin RN, Murray KO, Tapia-Conyer R, Gubler DJ: **Approaches to refining estimates of global burden and economics of dengue.** *PLoS Negl Trop Dis* 2014, **8**:e3306.